## IN THE CLAIMS

1. (CURRENTLY AMENDED) A photochromic diarylethene compound having isoxazole group expressed in the following formula (1),

$$\mathbb{R}^3$$
 $\mathbb{R}^3$ 
 $\mathbb{R}^2$ 
 $\mathbb{R}^2$ 

wherein  $R^1$  is a direct bond, O, or  $C_1$ - $C_3$  alkylene optionally substituted with-fluoro;  $R^2$  is a hydrogen atom,  $(CR^4H)_nOH$  or  $C_6(R5)_mH_1$ ;  $R^3$  is selected from the group consisting of a hydrogen atom, phenylisoxazole, hydroxymethylisoxazole, acetyl, hydroxy, and phenyl;  $R^4$  is hydrogen or  $C_1$ - $C_{10}$  alkyl;  $R^5$  is chloro, nitro, bromo, or the same as  $R^4$ ; X and Y are independently O, N, or S; Z is methylene optionally substituted with fluoro or carbonyl; and n, m and l are an integer of l to l.

- 2. (CURRENTLY AMENDED) A method for preparing the diarylethene compound of claim 1, comprising the steps:
- (i) formylating diarylethene compound of formula (2);
- (ii) reacting the formylated compound with NH<sub>2</sub>0H·HCl and aqueous basic solution in series and reacting with N-chlorosuccinimide (NCS); and
- (iii) reacting with acetylene compound substituted with R<sup>2</sup> in the presence of base catalyst,

$$\mathbb{R}^3$$
 $\mathbb{R}^1$ 
 $\mathbb{R}^1$ 
 $\mathbb{R}^1$ 
 $\mathbb{R}^3$ 
 $\mathbb{R}^3$ 
 $\mathbb{R}^3$ 

wherein  $R^1$  is a direct bond, O or  $C_1$ - $C_3$ -alkylene optionally substituted with fluoro;  $R^3$  is selected from the group consisting of a hydrogen atom, phenylisoxazole, hydroxymethylisoxazole, acetyl, hydroxy, and phenyl; X and Y are independently O, N, or S; and Z is methylene optionally substituted with a fluoro atom or carbonyl.

- 3. (WITHDRAWN) A photochromic diarylethene composition comprising 0.1-90wt.% of the compound of claim 1, 10-89.9wt.% of one or more resin selected from the group consisting of polyolefin, polycarbonate, polymethylmethacrylate, polyester, polyvinyl alcohol, polyurethane, and polyimide, and 10-89.9wt.% of one or more solvent.
- 4. (WITHDRAWN) A photochromic diarylethene composition comprising 0.1-90wt.% of the compound of claim 1, 10-99.8wt.% of fluorinated diacrylate monomer of formula (3), 0-80wt.% of monomer or oligomer having unsaturated group, 0.1-10wt.% of initiator of polymerization selected from thermalpolymerization initiator or photopolymerization initiator, and 0-90wt.% of one or more solvent,

$$CH_2 - CF_2 - CH_2$$
(3)

wherein n is an integer of 0 to 10.

5. (WITHDRAWN) The photochromic diarylethene composition according to claim 3, wherein said solvent is selected from the group consisting of acetone, hexane, acetonitrile,  $C_1$ - $C_{10}$  alcohol, dimethylformamide, tetraalkoxysilane, trialkoxysiliane,

dialkoxysilane, sulfuric acid, hydrochloric acid, organic acid, dimethylsulfoxide, pyridine, N-methylpyrrolidinone (NMP), sulfolane, α- methylnaphthalene, methoxynaphthalene, chloronaphthalene, diphenylethane, ethylene glycol, quinoline, dichloromethane, dichlorotoluene, propylene carbonate, xylene, methyl ethyl ketone, chloroform, methlylene chloride, trichloroethane, trichloroethylene, tetrahydrofuran, 1,4-dioxane and water.

- 6. (WITHDRAWN) The photochromic diarylethene composition according to claim 4, wherein said organic solvent is selected from the group consisting of acetone, hexane, acetonitrile, C<sub>1</sub>-C<sub>10</sub> alcohol, dimethylformamide, tetraalkoxysilane, trialkoxysiliane, dialkoxysilane, sulfuric acid, hydrochloric acid, organic acid, dimethylsulfoxide, pyridine, N-methylpyrrolidinone (NMP), sulfolane, α-methylnaphthalene, methoxynaphthalene, chloronaphthalene, diphenylethane, ethylene glycol, quinoline, dichloromethane, dichlorotoluene, propylene carbonate, xylene, methyl ethyl ketone, chloroform, methlylene chloride, trichloroethane, trichloroethylene, tetrahydrofuran, 1,4-dioxane and water.
- 7. (WITHDRAWN) The photochromic diarylethene composition according to claim 4, wherein said unsaturated monomer is selected from the group consisting of methyl methacrylate, butyl methacrylate, styrene, and α-methylstyrene.
- 8. (WITHDRAWN) The photochromic diarylethene composition according to claim 4, wherein said polymerization initiator is selected from the group consisting of benzoyl peroxide, 2,2'-azobisisobutyronitrile, and bis(l,1-dimethylethyl)peroxide, 1-hydroxycyclohexyl phenyl ketone, benzophenone, 2-hydroxy-1-[4-hydroxyethoxy)phenyl]-2-methyl-propanone, 2,2-dimethoxy-2-phenylacetophenone, fluorinated diaryltitanocine, and 2,2-bis(hydroxymethyl)propionic acid.
- 9. (WITHDRAWN) A photochromic diarylethene thin film prepared by coating a photochromic diarylethene compound having isoxazole group expressed by formula (1),

$$\mathbb{R}^3$$
 $\mathbb{R}^1$ 
 $\mathbb{R}^2$ 
 $\mathbb{R}^2$ 
 $\mathbb{R}^2$ 

wherein R<sup>1</sup> is a direct bond, O, or C<sub>1</sub>-C<sub>3</sub> alkylene optionally substituted with fluoro; R<sup>2</sup> is a hydrogen atom, (CR<sup>4</sup>H)<sub>n</sub>OH or C<sub>6</sub>(R5)<sub>m</sub>H<sub>1</sub>; R<sup>3</sup> is selected from the group consisting of a hydrogen atom, phenylisoxazole, hydroxymethylisoxazole, acetyl, hydroxy, and phenyl; R<sup>4</sup> is C<sub>1</sub>-C<sub>10</sub> alkyl; R<sup>5</sup> is chloro, nitro, bromo, or the same as R<sup>4</sup>; X and Y are independently O, N, or S; Z is methylene optionally substituted with fluoro or carbonyl; and n, m and l are an integer of 1 to 5, or and a photochromic thin film composition selected from the compositions of any of claims 2-6 on a substrate selected from conducting electrode substrate such as aluminum foil, aluminum drum, aluminum plate, platinum, Myler film, copper plate, conducting glass and conducting plastic; or a substrate such as polypropylene, propylene carbonate, polymethylmethacrylate, polyurethane, plastic, and glass.

10. (WITHDRAWN) A recording material, photochromic window, indicating element, plastic mirror, photochromic filter, photo switch, photosensitive drum, recording element, solar cell, lens, fiber, or optical element containing the compound selected from a photochromic diarylethene compound having isoxazole group expressed by formula (1),

$$\mathbb{R}^3$$
 $\mathbb{R}^1$ 
 $\mathbb{R}^2$ 
 $\mathbb{R}^2$ 

wherein  $R^1$  is a direct bond, O, or  $C_1$ - $C_3$  alkylene optionally substituted with fluoro;  $R^2$  is a hydrogen atom,  $(CR^4H)_nOH$  or  $C_6(R5)_mH_1$ ;  $R^3$  is selected from the group consisting of a hydrogen atom, phenylisoxazole, hydroxymethylisoxazole, acetyl, hydroxy, and phenyl;  $R^4$  is  $C_1$ - $C_{10}$  alkyl;  $R^5$  is chloro, nitro, bromo, or the same as  $R^4$ ; X and Y are independently O, N, or S; Z is methylene optionally substituted with fluoro or carbonyl; and n, m and l are an integer of 1 to 5, and one of said compositions of claims 2-6.

11. (NEW) A compound selected from the group consisting of 1-(6'-(5-hydroxymethylisoxazol)-2'-methylbenzo[b]thiophen-3'-yl)-2-(2"-methylbenzo[b]thiophen-3'-yl)hexafluorocyclopentene, 1-(6'-(5-phenylisoxazol)-2'-methylbenzo[b]thiophen-3'-yl)hexafluorocyclopentene, and di(6'-phenylisoxazol-2'-methylbenzo[b]thiophen-3'-yl)hexafluorocyclopentene.

## **CONCLUSION**

Wherefore, it is respectfully submitted that the claims are now in condition for allowance. Accordingly, it is respectfully requested that the claims be allowed and the case passed to issue.

Applicants' undersigned attorney may be reached in our New York office by telephone at (212) 218-2100. All correspondence should continue to be directed to our address given below.

Respectfully submitted,

Raymond R. Mandra
Attorney for Applicants
Registration No. 34,382

FITZPATRICK, CELLA, HARPER & SCINTO 30 Rockefeller Plaza
New York, New York 10112-3801
Facsimile: (212) 218-2200

NY\_MAIN 429468v2

w · ·